ANT/GRA/04

ARCTICA

GRANGE

R.C.M. 1970

0011

BOOK 2.

Loc. 247 —
351

INDEX

Page	Subject
1 – 13	~~Bosey Peaks~~ Mt. Sexton Area
14 – 18	Rookery Is. age det.
19 – 22	Mt. Burnett age det.
23 – 32	Scullin Monolith area
33 – 34.	Depot Peak area

N.B. all bearings magnetic.

SAT

4563/1627

299 On hill the dominant float is
a and py. gt's [illegible] granitic gneiss ± granit.
Also there are [illegible] py. granulite, mafic

~~gneiss~~ south 4364/1629

300 [illegible]

[illegible]

301 - 306 Photo AMS 24 Run 4 - 6'/85

307 Proposed [illegible]
to be sampled [illegible]
Dark band: 4364/1625
[illegible]

[illegible] produced [illegible] of the
samples. But much rather numerous
samples 3rd. Minor quantities, shears.

4363/1924

Sample 501) Most unusual & well worth
[illegible] over the [illegible] not much
[illegible] granules [illegible] similar
to [illegible] [illegible] [illegible]
with [illegible] [illegible]
[illegible] [illegible]
[illegible] [illegible]
[illegible] [illegible]
placed. Sample 502 (& sample 242)
[illegible] [illegible] [illegible]
[illegible] [illegible] [illegible]
[illegible] [illegible] [illegible]

4364/625

~~Sample~~ ?

Sample 304 (or 279)

(308) Discard shaft from Hill 303 [4375/622]
and cap and Hill 303
Red and dark brown sandstone flake

1/100,000 SR 42-43/9 C BEAVER LAKE

SUNDAY 2nd FEB 1975

All points marked on Photo AHT 94 Run 4C (7183)

Traverse with fine sample above working the northern run overlooking Diamond Glacier and White Basin. Steep with felsite gneiss

WC 309 Felsite porphyry on ridge 435/1660

Fresh coarse grained acid granulite dominantly yellowish v minor pyrite Glass-grey 45° Red-brown feldspars. Rare weathered red garnets Sample 309

Poor foliation due to [illegible] of feldspars 150° [illegible] sample site

Strongly jointed, blocky, craggy s/c block 1-2m joints 70°, 0°

Contains bands and lenses of streaky-weathered dark fine-grained biotite gneiss 30% mafics, some weathering. Contains [illegible] thick [illegible] Sample 310

Coarse!

Minor quartz veining [illegible] 030°

Various varieties of 309 due to [illegible] differences in mafics. Some peg. veining

[illegible]

7

[illegible] fine grained leucocratic veins
cutting the [illegible] rock from distance. They appear
to be more folding & irregularities in the foliation
[illegible] similar to those of Loc 309 & 302

same coordinates as for 309

Loc. 311 50° along ridge to east variants of 310
become dominant and form a migmatitic complex
with veins [illegible] felsic rock. Also diffuse
patches and lenses of [illegible] material. Most veins
are 0.5m or less wide and are approx. conformable
but diffuse patches of acid material are irregular
with the streaky mesocratic gneisses

Folds

Several banding structures and small similar
folds. Many veins have mafic schlieren. Usually
veins cross foliation to pegmatitic

[illegible] 350 / 50-60

Sample 311 of coarse grained granitic material
from one of veins

[illegible] fine to medium grained [illegible] mafic
[illegible] selvedge

Some development of felt [illegible] and augen gneiss
variants of 310.

[illegible] 8

+355/1663

Loc 312 Pebbly pelite ridge is composed of grey and light and weathering quartzite grain and grained Sample 312. A few pegmatitic bands composite ~ 5-10 cm thick Streaks of more mafic grain and a few minor bands less than 1 cm thick of pegmatite. Some development of fold pegmatite up to 5 cm Massive o/c

Fol 155 Banding 25°/90°

Nearby the quartzite grain size varies up to 10 cm thick of fine grained pegmatite. Sample (30% mafic) diffuse margins

But minor development of streaks of mafic with thickness 1" margins Sample 313

Major strike direction is 125°

Weathers grey green colour. Fresh is blue grey green slightly sandy

At least 3 cm thick vein dike [illegible] Appears to duplex some of banding with the granitic grain

Locality 12



as for 312

Loc 314 Upridge near summit similar rock type to 312 also granulite at 309 [illegible]
Flow development of segregations, internal and many angular fragments and veinings of
[illegible] segregations
Original development from rocks at loc 312.
Boundary - fine veins of 313 also cutting
[illegible]

Also at loc 314 [illegible] Sample 314
[illegible]

Descend scree to gully slope on E side of ridge. Traverse [illegible]
variety of granitic [illegible]
Main rock granite [illegible]

4368/1609

Near N end of ridge [illegible] high point
(loc 315) Massive ~~gl~~ pink feldspar granite
grain med. coarse [illegible] simple equigranular
up to 3cm small feld phenocrysts 1cm

11

Poorly foliated by streaks of muscovite alternating with felsic bands. Granoblastic. Accessory minerals are garnet, zircon, and opaque minerals. Matrix of quartz and feldspar grains. Quartz grains show undulose extinction. Feldspar grains show polysynthetic twinning and some are altered to sericite. Plagioclase shows normal zoning. Garnet porphyroblasts are fractured and contain inclusions of quartz and opaque minerals. Sericite alteration is present in the matrix.

Sample 315 (Gt-bi) Fine-grained granoblastic gneiss with garnet porphyroblasts. 4373/1672

(316) Fine-grained granoblastic gneiss with garnet porphyroblasts and abundant biotite. Biotite defines a foliation. Quartz and feldspar grains are equigranular. Garnet porphyroblasts are fractured and contain inclusions of quartz and opaque minerals. Sericite alteration is present in the matrix.

Sample 316. Fine-grained granoblastic gneiss with garnet porphyroblasts and abundant biotite. Biotite defines a foliation. Quartz and feldspar grains are equigranular. Garnet porphyroblasts are fractured and contain inclusions of quartz and opaque minerals. Sericite alteration is present in the matrix.

Fine-grained granoblastic gneiss with garnet porphyroblasts and abundant biotite. Biotite defines a foliation. Quartz and feldspar grains are equigranular. Garnet porphyroblasts are fractured and contain inclusions of quartz and opaque minerals. Sericite alteration is present in the matrix. (Sample 316)

Pebble

(317) 4373/1672 4378/1675

Fine-grained granoblastic gneiss with garnet porphyroblasts and abundant biotite. Biotite defines a foliation. Quartz and feldspar grains are equigranular. Garnet porphyroblasts are fractured and contain inclusions of quartz and opaque minerals. Sericite alteration is present in the matrix.

50-60° to margin. [redacted] fractured - blocks
to margin. Sample No. 317, similar to 279

[S] 330° 40°. Strike 310°, 50° to bearing
Strike parallel margin blocky for c/c
blocks about 2m. Several large channel
stones

Photos

(38)

WED 25th FEBRUARY Thurm SQ40-41/15

Rookery

All samples 4788/5001

Hanging

Sample 318 - Strong [illegible] Magn. shows [illegible] 50-55°. [illegible] dark [illegible] Massive o/c. Dark at 50° [illegible] 50° flat [illegible] Hanging [illegible] [illegible] 5m [illegible] Drill In U sample [illegible] [illegible] [illegible] 5 cm to 4m [illegible] [illegible] 55° [illegible] [illegible] [illegible] 35° [illegible]

[illegible] sample 318

Sample point about 15m S. from sample 317 222

Sample 319 Dom NNE for 318

Lith [illegible] [illegible] [illegible] to sample 318 Sheets a few cm [illegible] of [illegible] and [illegible] [illegible]. Tol. 35°

[illegible] [illegible] o/c [illegible] [illegible]

Sample [illegible] [illegible] [illegible]

[illegible] [illegible] [illegible] [illegible]

14


318
324
323

Sample 320 ✓ 10m. west from 321
Strata inclined towards ne 318°
310°. Strike pointed. 55°

[illegible] the [illegible] of [illegible]
[illegible] Cap [illegible] bands, [illegible]
[illegible] [illegible] [illegible]
310 [illegible] [illegible] [illegible]
[illegible] [illegible] [illegible]
[illegible] [illegible] [illegible]

[illegible] [illegible] [illegible]
[illegible] [illegible] [illegible]
[illegible] [illegible] [illegible]

1 the sample point.
Make a thin 10cm × 1.5m pre-
S.- ... sample pt. Use ground
geometric rather than ...
... to locate
... Blocky s/c

* NB. Do not refer to client samples 320

Sample 320 | 10m NW for 121

... ground ... ground
channels ... broken bedrock
Streaky foliation ... at 318, 24° ...
... strike ... somewhat more irregular
... 5m ... 3m ...
... 30° ... Approx 30° ... rock
Est 55-60°

... proof ... ground ... exposed
... (pit ... soil
...) ...
... the ...
... rich ... the ...
... 340° : 65° ... ...

grano blocks 2-3 m. diameter

Sample 323 *(Hammer sample)
12 m. ESE from 318
- tonalite Shirley, med.-coarse grained
Characteristic as 318.
No inclusions or mafic dominated areas
Shirley Lake stock - the largest mafic
[illegible] would contain [illegible] fine-
to coarse grained

Sample 324 *(Hammer sample)

[illegible] from 318, fine-coarse grained
Shirley characteristic as
318.

Between 323 & 324 several different
lithologies [illegible] - pegmatite
[illegible] - [illegible]

granit rock. Gt. pegs. ~ 2 cm
Vein ~ 10 cm wide, several in
a [illegible]. Also granular Lemon
quartz and [illegible] stringers and
veinlets

Sample 325 Grab sample from 318
Blanket [illegible]

Upper vein, Strike from 318
similar strong chalcocite coating
small vugs and in fractures, 1
pod [illegible] rock is less than 50%
pyrite, field graphs ~ 1 cm esp. in the
pods [illegible]

Large [illegible] strike and X cutting
[illegible] Glass. [illegible]
to the two main [illegible]

THURSDAY 26th FEB 1970

Mawson SQ 40-41/15

Mt. Burnett area. Age det. sample

about 100 m about ridge south from col S of Mt Burnett.

Sample 326 326-330 : 4595/4690

coarse grained Charnockite. Slightly porphyroblastic feldspars up to 2 cms. Differs from the Rookery Island - ever gneissic charnockite but not similar to the Mawson porph. charnockite. Orthoclase dominant felds. More granitic in texture & comp. ie feldspars are usually white or pale red/brown cf. Mawson purple feldspars.

Does not appear to be any alignment of feldspars or any gneissic texture

Minimal O/C.

Major joints 155°. Subsid joints at about 55°. Lineations marked.

Inclusions NN common near 326 but only small 20 cm x 10 cm lenses. Trend approx 60°

2 m for sample pit. Diffuse boundaries


N
328
6m
327

5m. east of sample pt area of large feld porph 3-4 cm long elongated ~ E-W plunge 60° over area of 2-3 m. = 5%

Occasional diffuse pinkish segregations of feldspar elongated ~ plunge 30°E

Minor anastomosing shears black, 11 irregular joints and healing joints. None with significant throw of 326. Shears dip shallowly S.

Thin ~ 2mm shear 2m to west of sample pt trend E-W; dip ~ 26°

White vein strike 215° 6-8 cm wide extends for ~ 15 m. Coarse 2 cm harrowed by east of sample pt. Diffuse ~ 1m zone with chloritic feld. porphyritic

Sample 326 V. fractured zone minor fe staining of

Samples 327, 328, 329 all similar massive coarse grained chloritic some feld. porphyry ⇒ 2cm

Sample 329 Minor fe weathering of feld

Sample 326 Weathering also fractures White deposit ⇒ fractures NSW v. good sample

327. Unsuccessful blasting. No sample


N
330
clos
331
Trunk

Sample location 330

Approx 75m north along ridge from 329 · 328
Fine-med grained charnockite forms an intrusion striking 265° approx west. Rather diffuse margin and no xenoliths.
Intrusion fine grained adj. to the country rock med-coarse grained charnockite some feld porph → 2cms. More mafics than at location 326 etc.
Mafic jointing in intrusion ~~some jointing at~~ 150° / intrusion at 30°
Country rock mafic joints 57° × 150°
Intrusion about 6m wide

Sample 331 Country rock 1-3m to SE of ~~qz~~ dyke.
Charnockite med-coarse grained. More mafic than 326 etc. Some irregular veins and pods of quartz material also some mafic segregations
Minor amounts of purple feld.

Small black amphibolite dikes similar to at 326 etc and similar style but none in immediate vicinity of sample points

NB Major joint approx vertical but most slabs are dipping at moderate angles towards the south

Sample 330. Kyle sandstone

Sample 331 Broke along the clear joint planes a few inches apart. Fe staining along joints and some white chemical staining.

Sample 332. Channel to [illegible] of col at [illegible] ridge. 4894/4095

3 miles $\frac{15}{4}$

FRIDAY 27th FEB 1970

Scullin Monolith SQ 42-43/13

Neill Peak

Locality (333) Mtk comprised N/S/W - base
Junction SW from Scullin Mount ll
Photo: ANT76 R.305 9031 R Grid Ref: 4004/4754
at northern end. Well layered rocks.
Main rock types are varieties of garnet
and granulite and some amounts of biotite
minor and very minor hornblende.
Samples 333 - 334. med. coarse grained.
Bands 10's of metres thick. thin layers of
marble ~ 1-2 m thick
Dark bands with the pink garnet
light coloured medium grained and
granulites etc are biotite gneisses often
rich in biotite sample 335 MMC
in garnet.
Sample 336 is a variety of the biotite gneiss
as above, several bands of pink garnet
basic material 1-2 m wide? Above,
sample 337 Another variety of the

23


336
333 c4
10-
337

dark bands contains phenocrysts of garnet > feld > qtz at less than 1 cm. Also shows appearance from bands up to 1 cm thick and not 336

Dark bands are up to 4 cm thick and cyclic form.

Sample 338 is garnet, acid granulite [illegible] and the comp. bands and also has angular garnet fragments 5 cm thick [illegible] angular fragments [illegible] up to 1 cm thick

Within each [illegible] there are minor irregular folds and deformation and the major bands and layers are [illegible] folding folded

General strike N40 E/145 Major fold 50°→225°

Irregular folds in minor gn fold [illegible] [illegible] of slumping

Folding in the layering appears to have been refolded - several domes of amphibole 4-5m seen on one ~~hand~~ surface
from the air
Knee-jerk ground the folding appears to have been tilted towards the top and along strike
One lineation trend observed 300 10° towards 300°
Mineralogical lineation
lineation itself appears to be folded along axes plunging 12° towards 250°

Sample 339. Thinned garnet acid granulite rock. Thinning has produced banding

Southern part of the is basic well banded mostly light coloured rock with dark folded veins

loc 340.

Scullin Monolith SQ42-43/13 · 4029/4787

Located near W end of Scullin pass.

Photo ANT41 R.17. 7070L

Migmatite and granite gneiss complex

Migmatite area. Dark mafic biotite gneiss and grained granulite with bands and pegmatite folds of pk feld ± mafics 1-2cm wide up to 10cm Sample 340

Bands leucos and irregular streaks of acid material

Also large irregular sp. to veins and masses of pk feld granulite

Sample 341 med to coarse grained

Extremely irregular and discontinuous association of 340 - 341 and other varieties eg 342 pk. granite gneiss + pegmatite

Cleavage 345 · 50° · 260°

Large gully → S-W side strikes 50°, contains float of Sample 343 pk. leuco gneiss and granite

to the marine + platy.

Composition of light + dark mineral
flecks in granitic gneiss - may be undigested

Several amphibolite veins (thin ones)
Foliation 320°

Sample 344 Variety of the mafic
granitic gneiss. contains felds. amphibolite
flecks parallel. Porph → dark Streaks,
due to mica banding.

Foliation irregular but about 240° steep dip
to west

Open folding of banding axis of which
here appear 250° vert.

To west of Sullom Voe main rock type is
coarse grained streaky granitic gneiss
The mafic material is flecky + patchy
displaced by the swirling of veins and
irregular masses of the granitic gneiss

until new landing spot there are only very occasional inclusions up to several inches diam. of mafic bi. gneiss. The actual relationship of the granite gneiss to the migmatite complex is difficult to determine because of vegetation, talus, etc.

Numerous measurements of shear strike at various exposures mainly 340° - 100° (three measurements per station) up to 30° west strike 100°.

The granite gneiss has some feldspar phenocrysts. Mafics less than 30%

loc 345 Sample 345

Sample 346 Red-grained variety occurring as irregular masses within 345 not to be confused

Strike - gneissosity 080° - 260° west

Min. lineation 30° towards 080°

Strike of joints 350° - 170°

Sample 347 Porphyroblastic coarse variety grained

Contains blocks and clasts → 1cm
wide x 20cm long of mafic granulite
showing minor pegmatitic veins in migmatites
and felsic gneisses

From air

Some ridges are seen with
X-cutting gullies with possible dykes
Extend across whole mountain.

1 rock outcrop of major peak but did
rocks appear similar. Is there at
333 etc.

N peak is main is rock granulite with
minor dykes ~~back~~ is. Undeformed pegmatite o/c

Western peak of Mount Monteith
Loc 348 ANT 41. R.17. 8066 Smith Kop SQ42-43/13
4103/4894
Coarse granoid gt-opx-qtz-felsic hornfels
from Loc 348 → 4 cm

Sample 348

Thermo-deformed o/c
Strikes 65° - 125°
Several ~~gt~~ pegs veins → 10 cm wide
in length of 50

[illegible] = fine micaceous sandstone and sandy [illegible] fine metres of micaceous [illegible]

Also occasional very [illegible] with [illegible] and micaceous [illegible] [illegible]

[illegible] foliation planes mainly 70° – 80° also N/S [illegible]

Outcrop 405m. N of Mervyn Monolith

Loc 349 ANT. 41. R17. 7067L. 4091/4779

Banded micaceous fine-grained sandstone. Alternating bands of micaceous and [illegible] 1–2 cm and up to 10 cm.

Sample 349 Similar to sample 342

Irregular pods [illegible] and bands of medium-grained pinkish-brown weathering rock. Sample 350

Also veining veins → 1–2 cm wide of [illegible] 350

Bedding: 190/15 [illegible]

Foliation planes 10° [illegible]

Enough

N face.

Enough outcrop to find dip and all
dykes folded fault. Open anticline
upto 10m

Drill a few bores.

NB Note difference in lithology between
borehole 348 & 349

Drill 349 in syncline to a depth of
Normal Permian 349 in the middle borehole
should intersect Coal Measures
at a depth of 150m
Bands of [illegible] 4-5m thick
Several [illegible] upto [illegible]
veins (350) will also now [illegible]

themes and terms

On west side of hill a pink coloured
aastvein + few metres wide was observed
X altig face at an angle



SUNDAY 1st MARCH DEPOT PEAK AREA

Loc 351 Stinnes Nunks. SR 40-41/85.

Small low nunk situated about 5 miles & 5 km
about 150-160° from Depot Peak
Mag. bearing of 50° from o/c to Depot Peak
O/c about 100 yds long elongated about
300° mag.
Rock fluted on sides and rubble except at
NE
northern end where o/c is intact
Varieties of qtz bi qtz felsite granite. Basalt
felsite Standing stone Conglomerate vesicular
Samples 351 - 352
Med. coarse grained
Felsite porph. 1 up to 1½ cms ?

Felsite 200 x 15 Zone 41: 5680/3333

A few flakes of black vitreous material

Sample 353 shows weathering to brown
Weathers brown/yellow with qtz standing out
as blue grey
Acid granulite with biotite. Mineral banding

due to segregation of 93° feldspars into layers 1–2 mm wide.

10000
2360
12360

1:100,000

1 mile = .06 inches
1 inch = 13 miles



10000
2536
30,540

1:100,000

1 mile = 0.6 inches.
1 inch = 1.3 miles